

On Automorphisms of Matrix Invariants Induced From the Trace Ring

Zinovy Reichstein
Department of Mathematics
University of California
Berkeley, California 94720

Submitted by Robert M. Guralnick

ABSTRACT

1. INTRODUCTION

In this paper we continue the study of the automorphism group of the space $Q_{m,n}$ initiated in [14]. Recall that $Q_{m,n}$ is the algebraic quotient space for the simultaneous conjugation action of $\mathrm{PGL}_n(k)$ on the affine space $M_n(k)^m$ of m -tuples of $n \times n$ matrices. Here and throughout the paper the base field k will be assumed to be algebraically closed and of characteristic 0.

Recall that the affine variety $Q_{m,n}$ is stratified according to representation type. The representation type of a point $p \in Q_{m,n}$ is defined as follows. Let $(X_1, \dots, X_m) \in M_n(k)$ be an element of the unique closed PGL_n orbit which projects to p under the quotient map. Then the representation

$$\begin{aligned} \phi : k\{u_1, \dots, u_m\} &\rightarrow M_n(k), \\ u_i &\rightarrow X_i \end{aligned} \tag{1}$$

of the free algebra $F_m = k\{u_1, \dots, u_m\}$ is semisimple; see [1]. We say that p has representation type $\tau = (e_1, d_1; \dots; e_r, d_r)$ if ϕ is the sum of r irreducible representations ϕ_i of dimension d_i and multiplicity e_i . The set of all points of representation type τ will be denoted by $Q_{m,n}(\tau)$. The group of algebraic representation-type-preserving automorphisms of $Q_{m,n}$ will be denoted by $\mathbf{Aut}(Q_{m,n})$.

In [14] we proved the following result.

THEOREM 1.1. *Assume $m \geq n + 1$. Then*

(a) [14, Theorem 1.2] $\mathbf{Aut}(Q_{m,n})$ acts transitively on $Q_{m,n}(\tau)$ for any representation type τ .

(b) [14, Theorem 1.3] $\mathbf{Aut}(Q_{m,n})$ acts s -transitively on $Q_{m,n}(1, n)$ for any positive integer s .

The automorphisms of $Q_{m,n}$ which appear in the proof of Theorem 1.1 are all induced by automorphisms of the free polynomial algebra $F_m = k\{u_1, \dots, u_m\}$. In this paper we study the broader class of automorphisms of $Q_{m,n}$ induced by k -algebra automorphisms of the trace ring $T_{m,n}$. For generalities on such automorphisms see Section 3. These new automorphisms allows us to strengthen Theorem 1.1 as follows.

THEOREM 1.2. *Assume $m \geq n + 1$. Then for any positive integer s the group $\mathbf{Aut}(Q_{m,n})$ acts s -transitively on the stratified space $Q_{m,n}$. In other words, suppose (p_1, \dots, p_s) and (q_1, \dots, q_s) are two s -tuples of distinct points in $Q_{m,n}$ such that $\tau(p_i) = \tau(q_i)$ for $i = 1, \dots, s$. Then there exists a $g \in \mathbf{Aut}(Q_{m,n})$ such that $g(p_i) = q_i$ for $i = 1, \dots, s$.*

We give a proof of this theorem in Section 5. The condition $m \geq n + 1$ is necessary in the case $n = 1$, since every automorphism of $Q_{1,1} = k^1$ is of the form $x \rightarrow ax + b$. We do not know whether or not the theorem holds for $m \leq n$ when $n \geq 2$. We also note that for some choices of $\{p_i\}, \{q_i\}$ the element g in the statement of Theorem 1.2 cannot be induced by an automorphism of the polynomial algebra F_m ; see Remark 3.4.

As we mentioned above, an automorphism of F_m induces an automorphism of $T_{m,n}$ which in turn induces a representation type preserving automorphism of $Q_{m,n}$; see Section 3 for details. In Sections 6 and 7 we study the resulting diagram of group homomorphisms

$$\mathrm{Aut}(F_m) \rightarrow \mathrm{Aut}(T_{m,n}) \rightarrow \mathbf{Aut}(Q_{m,n}).$$

The main result of these sections is that the second map is not surjective. In other words,

THEOREM 1.3. *Suppose $n \geq 3$ and $m \geq 2$. Then there exists a $g \in \mathbf{Aut}(Q_{m,n})$ which is not induced by an automorphism of the trace ring $T_{m,n}$.*

The automorphism g in the theorem is, in fact, of a very simple form; see Proposition 7.2. The key part of our proof of Theorem 1.3 is the criterion for $g \in \mathrm{Aut}(T_{m,n})$ to be trivial given by Corollary 6.2. This criterion is closely related to a theorem of Guralnick and Montgomery [5]. In Section 6 we present a proof of this theorem based on our criterion.

2. PRELIMINARIES

The Trace Ring $T_{m,n}$

Let $x_{ij}^{(l)}$ be mn^2 independent commuting variables. Here $i, j = 1, \dots, n$ and $l = 1, \dots, m$. The ring of generic matrices $G_{m,n}$ is the k -subalgebra of the matrix algebra $M_n(k[x_{ij}^{(l)}])$ generated by the m elements $X_1 = (x_{ij}^{(1)}), \dots, X_m = (x_{ij}^{(m)})$. The trace ring $T_{m,n}$ is defined as the subalgebra of $M_n(k[x_{ij}^{(l)}])$ generated by elements of $G_{m,n}$ and the coefficients of their characteristic polynomials. The latter are viewed as scalar matrices in $M_n(k[x_{ij}^{(l)}])$. In fact, since our base field k is assumed to be of characteristic 0, Newton's formulas say that $T_{m,n}$ is generated by elements of $G_{m,n}$ and their traces. Denote the subalgebra of scalar elements of $T_{m,n}$ by $C_{m,n}$, i.e.

$$C_{m,n} = k[x_{ij}^{(l)}] I_{n \times n} \cap T_{m,n}.$$

In other words, $C_{m,n}$ is the k -algebra generated by elements of the form $\text{tr } Z$ for $Z \in G_{m,n}$.

Note that the natural $(Z^+)^m$ -grading of $G_{m,n}$ by the multidegree in (X_1, \dots, X_m) extends to a $(Z^+)^m$ -grading of $T_{m,n}$. We shall denote the degree of an element $Z \in T_{m,n}$ with respect to X_i by $\deg_i Z$ and the total degree of Z by $\deg Z$.

The Universal Property of $T_{m,n}$

As a direct consequence of the above definition, the trace ring $T_{m,n}$ has the following universal property.

PROPOSITION 2.1. *Let R be a commutative k -algebra, N be a subalgebra of the matrix algebra $M_n(R)$ such that $\text{tr } N \subset N$, and $a_1, \dots, a_m \in N$. Then there exists a unique k -algebra homomorphism $h: T_{m,n} \rightarrow N$ such that $h \circ \text{tr} = \text{tr} \circ h$ and $h(X_l) = a_l$ for $l = 1, \dots, m$.*

Indeed, we obtain h by specializing each $x_{ij}^{(l)}$ to the (i, j) entry of a_l .

We remark that one can prove a similar statement about any algebra N with trace by using the embedding theorem of Procesi [12]; see [14, Lemma 2.8]. However, in the sequel we will only appeal to the elementary version of this result stated above. As a corollary we obtain the following fact.

COROLLARY 2.2. *The natural embedding $G_{m-1,n} \hookrightarrow G_{m,n}$ which takes X_l to X_l for $l = 1, \dots, m-1$ extends to an embedding $T_{m-1,n} \hookrightarrow T_{m,n}$.*

Proof. Applying Proposition 2.1 to the $m-1$ -tuple $X_1, \dots, X_{m-1} \in T_{m,n}$, we obtain a homomorphism $h: T_{m-1,n} \rightarrow T_{m,n}$. To see that h is

injective, we apply Proposition 2.1 once again. It says that there is a homomorphism $g: T_{m,n} \rightarrow T_{m-1,n}$ such that $g(X_l) = X_l$ for $l = 1, \dots, m-1$ and $g(X_m) = 0$. Then $g \circ h(X_l) = X_l$. Moreover, $g \circ h: T_{m-1,n} \rightarrow T_{m-1,n}$ commutes with the trace function. Hence, $g \circ h$ is the identity map on $T_{m-1,n}$. In particular, h is injective. ■

REMARK 2.3. In the sequel we shall identify $T_{m-1,n}$ with its image in $T_{m,n}$.

Properties of $T_{m,n}$ and $C_{m,n}$

We shall make use of the following properties of $T_{m,n}$ and $C_{m,n}$.

PROPOSITION 2.4. *Any nonzero element $Z \in T_{m,n}$ is nonsingular, i.e., $\det Z \neq 0$. In particular, $T_{m,n}$ is a domain.*

This is a consequence of the fact that $C_{m,n}$ is a domain and $T_{m,n}$ embeds in its total field of fractions, which is a division algebra. See [2, Section 12.6] for details.

PROPOSITION 2.5 (Procesi [11]). *$C_{m,n}$ is the center of $T_{m,n}$.*

PROPOSITION 2.6 (Sibirskii [16], Procesi [11]). *$C_{m,n}$ is the ring of invariants for the conjugation action of $\mathrm{PGL}_n(k)$ on $M_n(k)^m$. In other words, $C_{m,n}$ is the coordinate ring of the affine variety $Q_{m,n}$.*

PROPOSITION 2.7 (Shirshov [15, Section 4.2]). *$T_{m,n}$ is finitely generated as $C_{m,n}$ -module.*

We refer the reader to [4], [6], and [13] for a more detailed account of the properties of the trace ring.

Automorphisms of $T_{m,n}$ Commute with tr

In the sequel the characteristic polynomial of a matrix M will be denoted by

$$\mathrm{char}_M(t) = t^n + \sum_{i=0}^{n-1} (-1)^i c_i(M) t^{n-i}.$$

The group of k -algebra automorphisms of $T_{m,n}$ will be denoted by $\mathrm{Aut}(T_{m,n})$

PROPOSITION 2.8. *Let f be a k -algebra automorphism of $T_{m,n}$. Then for any $Z \in T_{m,n}$ and any $i = 0, 1, \dots, n-1$ we have $c_i(f(Z)) = f(c_i(Z))$. In particular, $\mathrm{tr} f(Z) = f(\mathrm{tr} Z)$ and $\det f(Z) = f(\det Z)$.*

Proof. By the Cayley-Hamilton identity an element $Z \in T$ satisfies two monic polynomials of degree n :

$$\text{char}_Z(t) = t^n + \sum_{i=0}^{n-1} (-1)^i c_i(Z) t^{n-i},$$

$$f^{-1} \text{char}_{f(Z)}(t) = t^n + \sum_{i=0}^{n-1} (-1)^i f^{-1}[c_i(f(Z))] t^{n-i}.$$

If Z has distinct eigenvalues, then these polynomials must coincide. That is, $c_i(Z) = f^{-1}c_i(f(Z))$, as desired. Now let Z be an arbitrary element of $T_{m,n}$. Note that X_1 has distinct eigenvalues, and hence so does $Z_s = Z + sX_1$ for all but finitely many $s \in k$. Then $c_i(Z_s) = f^{-1}(c_i(f(Z_s)))$ holds for all but finitely many $s \in k$; hence, it holds for all $s \in k$, and in particular, for $s = 0$. ■

COROLLARY 2.9. *Let f be a k -algebra automorphism of $T_{m,n}$. Then f is trivial if and only if $f(X_i) = X_i$ for $i = 1, \dots, m$.*

Proof. Suppose $f(X_i) = X_i$ for $i = 1, \dots, m$. Then $f(Z) = Z$ for every $Z \in G_{m,n}$. By Proposition 2.8 $f(c_i(Z)) = c_i(f(Z)) = c_i(Z)$ for every $Z \in G_{m,n}$ and every $i = 0, 1, \dots, n-1$. Since $T_{m,n}$ is generated by elements of the form Z and $c_i(Z)$ as Z ranges over $G_{m,n}$, the automorphism f is trivial on all of $T_{m,n}$. ■

PROPOSITION 2.10. *Let Λ be the collection of all m -tuples $Y_1, \dots, Y_m \in T_{m,n}$ which generate $T_{m,n}$ as a k -algebra with trace. Then the correspondence*

$$\text{Aut}(T_{m,n}) \rightarrow \Lambda,$$

$$f \rightarrow (f(X_1), \dots, f(X_m))$$

is a bijection.

Proof. Injectivity follows from Corollary 2.9. To prove surjectivity suppose Y_1, \dots, Y_m generate $T_{m,n}$ as an algebra with trace. Then by Proposition 2.1 there exists an endomorphism $f: T_{m,n} \rightarrow T_{m,n}$ which takes X_1, \dots, X_m to Y_1, \dots, Y_m respectively. Since f commutes with tr , it is surjective. It

remains to show that f is also injective. Assume the contrary. Then

$$(0) \subset \text{Ker } f \subset \text{Ker } f^2 \subset \dots$$

is a strictly increasing sequence of $C_{m,n}$ -submodules of $T_{m,n}$ contradicting Proposition 2.7. \blacksquare

Representations

Let M be a k -algebra. A *representation* (an *irreducible representation*) of F_m in M is a k -algebra homomorphism (a surjective k -algebra homomorphism)

$$\rho : F_m \rightarrow M.$$

Two representations $\rho_1 : F_m \rightarrow M_1$ and $\rho_2 : F_m \rightarrow M_2$ are said to be *equivalent* if there exists an isomorphism $f : M_1 \rightarrow M_2$ such that $\rho_2 = f \circ \rho_1$. We write $\rho_1 \approx \rho_2$.

A representation ϕ is called *semisimple* if its image is a finite-dimensional semisimple k -algebra, i.e. a finite direct sum of matrix algebras over k . A semisimple representation can thus be viewed as a direct sum of irreducible representations in matrix algebras over k . Grouping the equivalent ones together, we write

$$\phi = \phi_1^{e_1} \oplus \dots \oplus \phi_r^{e_r},$$

where $\phi_i : F_m \rightarrow M_{d_i}(k)$ is irreducible for $i = 1, \dots, r$ and $\phi_i \not\approx \phi_j$ for $i \neq j$. As we mentioned in the introduction, every point $p \in Q_{m,n}$ is represented by an m -tuple of $n \times n$ matrices (X_1, \dots, X_m) such that the representation

$$\rho : F_m \rightarrow M_n(k),$$

$$u_l \rightarrow X_l$$

for $l = 1, \dots, m$ is semisimple. Up to equivalence ρ is unique; we shall refer to it as the semisimple representation associated to p . Suppose

$$\rho = \rho_1^{e_1} \oplus \dots \oplus \rho_r^{e_r}, \tag{2}$$

where $\rho_i : F_m \rightarrow M_{d_i}(k)$ is irreducible for $i = 1, \dots, r$ and $\rho_i \not\approx \rho_j$ for $i \neq j$.

Then we define the representation type of p (or ρ) to be the unordered r -tuple of pairs (e_j, d_j) and write

$$\tau(p) = (e_1, d_1; \dots; e_r, d_r).$$

Note that $\sum_{i=1}^r e_i d_i = n$.

If the semisimple representation associated to p is as in (2), then we define the reduced semisimple representation

$$\phi: F_m \rightarrow M_{d_1} \times \dots \times M_{d_r}$$

associated to p by $\phi = \rho_1 \oplus \dots \oplus \rho_r$.

PROPOSITION 2.11 [1, Theorem 9.2; 14, Lemma 2.4]. *For $i = 1, \dots, r$ let ϕ_i be representation of F_m in a matrix algebra over k . Then $\rho = \rho_1 \oplus \dots \oplus \rho_r$ is irreducible if and only if ρ_1, \dots, ρ_r are irreducible and pairwise inequivalent.*

In particular, the reduced semisimple representation associated to $p \in Q_{m,n}$ is always irreducible.

3. WELL-BEHAVED s -TUPLES

The automorphisms of $Q_{m,n}$ which will be used in our proof of Theorem 1.2 will all be induced from automorphisms of $T_{m,n}$ in the following simple way. Let f be a k -algebra automorphism of $T_{m,n}$. Since $C_{m,n}$ is the center of $T_{m,n}$ (Proposition 2.5), f induces an automorphism of $C_{m,n}$. We denote the dual automorphism of $Q_{m,n} = \text{Spec } C_{m,n}$ by f_* ; see Proposition 2.6.

It is easy to see that f_* can also be described as follows. Consider the automorphism f^* of $k^{mn} = M_n(k)^m$ (as an affine variety) given by

$$A = (A_1, \dots, A_m) \rightarrow (f(X_1)(A), \dots, f(X_m)(A)).$$

This automorphism commutes with the conjugation action of PGL_n on $M_n(k)^m$. Hence, it descends to an automorphism f_0 of the quotient variety $Q_{m,n}$; see [10, Theorem 1.1; 14, Lemma 2.2].

We claim that $f_* = f_0$. Indeed, suppose $p \in Q_{m,n}$ is represented by $(A_1, \dots, A_m) \in M_n(k)^m$. It is enough to check that $f(c)(p) = c(f_0(p))$ for every c in $C_{m,n}$. Both sides are k -algebra homomorphisms $C_{m,n} \rightarrow k$. Thus

we may assume without loss of generality that $c = \text{tr}(X_{i_1} \cdots X_{i_N})$ for some $1 \leq i_1, \dots, i_N \leq m$. Then by Proposition 2.8

$$f(c)(p) = \text{tr} f(X_{i_1} \cdots X_{i_N})(p) = \text{tr} f^*(A_{i_1} \cdots A_{i_N}) = c(f_0(p)),$$

as desired.

Let σ be an automorphism of the free algebra $F_m = k\{u_1, \dots, u_m\}$. For $j = 1, \dots, m$ let Y_j be the image of $\sigma(u_j)$ in $T_{m,n}$ under the map $F_m \rightarrow T_{m,n}$ which takes u_i to X_i . Then the k -algebra generated by Y_1, \dots, Y_m contains X_1, \dots, X_m . Hence, Y_1, \dots, Y_m generate $T_{m,n}$ as an algebra with trace. By Proposition 2.10 there is a unique k -algebra automorphism of $T_{m,n}$ which takes X_i to Y_i for $i = 1, \dots, m$. We denote this automorphism by σ' . Note that by our description of f_* above, σ'_* is exactly the automorphism of $Q_{m,n}$ which we denoted by σ_* in [14].

LEMMA 3.1. *Let f be a k -algebra automorphism of $T_{m,n}$. Then f_* is a representation-type-preserving automorphism of $Q_{m,n}$.*

Proof. The argument of the proof of [14, Proposition 4.1] goes through unchanged in this case. We briefly recall it here. Let p be a point of $Q_{m,n}$ represented by a semisimple m -tuple of matrices $A = (A_1, \dots, A_m)$. Then by the definition of partial order on the representation types we have $\tau(A) \geq \tau(f^*(A))$, i.e. $\tau(p) \geq \tau(f_*(p))$. Applying the same argument to f^{-1} , we obtain equality. ■

We thus have a sequence of group homomorphisms

$$\text{Aut}(F_m) \rightarrow \text{Aut}(T_{m,n}) \rightarrow \mathbf{Aut}(Q_{m,n})$$

where the first map is given by $\sigma \rightarrow \sigma'$ and the second map is given by $f \rightarrow f_*$. We shall return this sequence in Section 7.

We now turn our attention to Theorem 1.2.

DEFINITION 3.2. We call an s -tuple $p_1, \dots, p_s \in Q_{m,n}$ *well behaved* if no two of the associated semisimple representations $\rho_1, \dots, \rho_s : F_m \rightarrow M_n(k)$ have an isomorphic irreducible component.

PROPOSITION 3.3. *The assertion of Theorem 1.2 holds if (p_1, \dots, p_s) and (q_1, \dots, q_s) are well-behaved s -tuples. Moreover, in this case there is a $\sigma \in \text{Aut}(F_m)$ such that $\sigma'_*(p_i) = q_i$ for $i = 1, \dots, s$.*

Proof. Let $\rho_i, \mu_i : F_m \rightarrow M_n(k)$ be the semisimple representations associated to p_i and q_i respectively. Denote the representation type of p_i by $\tau = (e_{i1}, d_{i1}; \dots; e_{ir_i}, d_{ir_i})$; by our assumption τ is also the representation type of q_i . Write

$$\rho_i = \rho_{i1}^{e_{i1}} \oplus \dots \oplus \rho_{ir_i}^{e_{ir_i}} \quad \text{and} \quad \mu_i = \mu_{i1}^{e_{i1}} \oplus \dots \oplus \mu_{ir_i}^{e_{ir_i}}, \quad (3)$$

where $\rho_{ij}, \mu_{ij} : F_m \rightarrow M_{d_{ij}}(k)$ are irreducible representations and $\rho_{ij} \not\cong \rho_{il}, \mu_{ij} \not\cong \mu_{il}$ for $j \neq l$.

By our assumption the representations ρ_{ij} are pairwise inequivalent and the representations μ_{ij} are pairwise inequivalent. Hence, the representations

$$\rho = \bigoplus_{i,j} \rho_{ij} \quad \text{and} \quad \mu = \bigoplus_{ij} \mu_{ij} : F_m \rightarrow \bigoplus_{i,j} M_{d_{ij}}(k)$$

have the same representation type; namely $(1, d_{11}; \dots; 1, d_{sr_s})$. By Proposition 2.11 ρ and μ are irreducible. Since each $d_{ij} \leq n \leq m+1$, we can appeal to [14, Theorem 4.4], which says that $\rho_{ij} \circ \sigma = \mu_{ij}$ for some $\sigma \in \text{Aut}(F_m)$. Therefore, $\rho_i \circ \sigma = \mu_i$ and hence $\sigma'_*(p_i) = q_i$ for $i = 1, \dots, s$. ■

REMARK 3.4. Note that the condition that the s -tuples (p_1, \dots, p_s) and (q_1, \dots, q_s) are well behaved is essential here. Indeed, for any $\sigma \in \text{Aut}(F_m)$, the s -tuple $(\sigma_* p_1, \dots, \sigma_* p_s)$ is well behaved if and only if (p_1, \dots, p_s) is well behaved. This follows from the fact that representations $\rho, \mu : F_m \rightarrow M$ are equivalent if and only if $\rho \circ \sigma$ and $\mu \circ \sigma$ are equivalent. In particular, an element of $\mathbf{Aut}(Q_{m,n})$ of the form σ'_* cannot carry a well-behaved s -tuple into one which is not well behaved. As we shall see in Section 5, we can get around this difficulty by considering elements of $\mathbf{Aut}(Q_{m,n})$ of the form f_* for $f \in \text{Aut}(T_{m,n})$.

4. TAME AUTOMORPHISMS

In this section we discuss some properties of the group $\text{TAut}(T_{m,n})$ of tame automorphisms of $T_{m,n}$. We shall view $\text{TAut}(T_{m,n})$ as an infinite-dimensional algebraic group as defined by Shafarevich in [17]. We will show that this group is covered by an ascending collection $\{G_t\}$ of irreducible finite-dimensional varieties. For this reason, $\text{TAut}(T_{m,n})$ behaves in some instances as an irreducible linear algebraic group. The results of this section will be used in the proof of Theorem 1.2 in Section 5.

Let σ be an automorphism of the free polynomial algebra $F_m = k\{u_1, \dots, u_m\}$. Recall that σ is linear if it is given by

$$g(u_i) = \sum_{j=1} a_{ij} u_j,$$

where $a_{ij} \in k$ and the matrix (a_{ij}) is nonsingular. The automorphism σ is lower triangular if for every $i = 1, \dots, m$ there exists a polynomial P_i in $i - 1$ variables such that $\sigma(u_i) = u_i + P_i(u_1, \dots, u_{i-1})$.

Similarly, an automorphism f of $T_{m,n}$ is linear if

$$f(X_i) = \sum_{j=1} a_{ij} X_j$$

where $a_{ij} \in k$ and the matrix (a_{ij}) is nonsingular. An automorphism of $T_{m,n}$ is lower triangular if it is of the form

$$\begin{aligned} X_1 &\rightarrow X_1, \\ X_2 &\rightarrow Y_1 + X_2, \\ &\vdots \\ X_m &\rightarrow Y_{m-1} + X_m, \end{aligned} \tag{4}$$

where Y_i is an element of $T_{i,n}$. Here we view $T_{i,n}$ as a subring of $T_{m,n}$ for $i = 1, \dots, m - 1$; see Remark 2.3.

An automorphism of F_m [or $T_{m,n}$] is called *tame* if it can be written as a composition of linear and lower triangular automorphisms. Tame automorphisms form a subgroup of $\text{Aut}(F_m)[\text{Aut}(T_{m,n})]$, which is denoted by $\text{TAut}(F_m)$ [$\text{TAut}(T_{m,n})$]. Conjecturally $\text{TAut}(F_m) = \text{Aut}(F_m)$ and $\text{TAut}(T_{m,n}) = \text{Aut}(T_{m,n})$.

As a direct consequence of our definitions we obtain the following lemma.

LEMMA 4.1. *Let σ be a tame automorphism of F_m . Then σ' is a tame automorphism of $T_{m,n}$.*

Let $G(d) = \{h_1 g_1 \cdots h_d g_d\}$, where each h_j is linear and each g_i is lower triangular of degree $\leq d$. Here by the degree of an automorphism f of $T_{m,n}$ we mean the maximum of the total degrees of $g(X_l)$ as $l = 1, \dots, m$; see Section 2. Clearly $\text{TAut}(T_{m,n}) = \bigcup_{d \geq 1} G_d$.

Note that an element of G_d is of degree at most d^d . In other words, if Z_1, \dots, Z_N is a basis of the k -vector space of elements of $T_{m,n}$ of degree $\leq d^d$, then

$$f(X_i) = \sum_{j=1}^N c_{ij} Z_j \quad (5)$$

for some $c_{ij} \in k$. The scalars c_{ij} can be thought of as the coordinates of f . Since an element $f \in G_d$ is uniquely determined by its mN coordinates c_{ij} , we can think of G_d as a subset of the affine space k^{mN} .

LEMMA 4.2. G_d is an irreducible constructible subset of k^{mN} .

Proof. Let L_d be the set of lower triangular automorphisms of $T_{m,n}$ of degree $\leq d$. Since each Y_i in (4) ranges over the finite-dimensional k -vector space of elements of $T_{i,n}$ of degree $\leq d$, L_d is isomorphic to an affine space. Thus we have a regular surjective map

$$(\mathrm{GL}_m(k) \times L_d)^d \rightarrow G_d$$

given by $(h_1, g_1, \dots, h_d, g_d) \rightarrow h_1 g_1 \cdots h_d g_d$. Its image is constructible by Chevalley's theorem. Since $\mathrm{GL}_m(k) \times L_d$ is irreducible, so is G_d . ■

LEMMA 4.3. Let $p, q \in Q_{m,n}$, and let W be the set of all $f \in G_d$ such that $(f_* p, f_* q)$ is well behaved. Then W is Zariski-open in G_d .

Proof. Suppose the semisimple representations associated to p and q are

$$\rho = \rho_1^{a_1} \oplus \cdots \oplus \rho_r^{a_r} \quad \text{and} \quad \mu = \mu_1^{b_1} \oplus \cdots \oplus \mu_s^{b_s},$$

where $\rho_i, \mu_i : F_m \rightarrow M_{d_i}(k)$ are irreducible.

Let $f \in G_d$ be as in (5). Then the semisimple representations associated to $f_* p$ and $f_* q$ are of the form

$$\rho(c) = \rho_1(c)^{a_1} \oplus \cdots \oplus \rho_r(c)^{a_r} \quad \text{and} \quad \mu(c) = \mu_1(c)^{b_1} \oplus \cdots \oplus \mu_s(c)^{b_s},$$

where $\rho_i(c), \mu_i(c)$ are representations of F_m in $M_{d_i}(k)$ which depend on $c = (c_{ij})$.

Fixing p and q and letting f vary over G_d , we see that for every $P \in F_m$ the entries of $\rho_i(c)(P)$ and $\mu_i(c)(P)$ are polynomials in c_{ij} . Moreover, for $f \in G_d$ the representation type of $f_* p$ is the same as the representation type of p ; see Lemma 3.1. Hence, the representations $\rho_i(c): F_m \rightarrow M_{d_i}(k)$ are irreducible. Similarly each $\mu_i(c)$ is irreducible.

Therefore, $(f_* p, f_* q)$ is not well behaved iff for some $1 \leq \alpha \leq r$ and $1 \leq \beta \leq s$ the representations $\rho_\alpha(c)$ and $\mu_\beta(c)$ are equivalent. Since they are irreducible, this happens if and only if

$$\mathrm{tr} \rho_\alpha(c)(P) = \mathrm{tr} \mu_\beta(c)(P)$$

for every $P \in F_m$.

These are polynomial conditions on c_{ij} . They cut out a closed subset of G_d for every α and β . The set W is the complement of the union of these closed sets taken over all pairs (α, β) with $\alpha = 1, \dots, r$ and $\beta = 1, \dots, s$. Therefore, W is Zariski-open in G_d . ■

Recall that a representation ρ of F_m is *saturated* if its restriction to $F_{m-1} = k\{u_1, \dots, u_{m-1}\}$ is irreducible. A point $p \in Q_{m,n}$ is saturated if its reduced semisimple representation is saturated. In other words, p is saturated if its associated semisimple representation ρ has the property that $\rho(F_{m-1}) = \rho(F_m)$.

LEMMA 4.4. *Let $p \in Q_{m,n}$. Then the set*

$$\{f \in G_d : f_* p \text{ is saturated}\}$$

is Zariski-open in G_d .

Proof. Denote the semisimple representations associated to p by

$$\rho = \rho_1^{e_1} \oplus \dots \oplus \rho_r^{e_r},$$

where $\rho_i: F_m \rightarrow M_{d_i}(k)$ are inequivalent irreducible representations.

Let $f \in G_d$ be as in (5). As we saw in the proof of the previous lemma, the semisimple representation associated to $f_* p$ is of the form

$$\rho(c) = \rho_1(c)^{e_1} \oplus \dots \oplus \rho_r(c)^{e_r}.$$

Here the representations $\rho_1(c), \dots, \rho_r(c)$ are inequivalent and irreducible, and the entries of the matrix $\rho_i(c)(P)$ are polynomials in c_{ij} for every

$P \in F_m$ and every $i = 1, \dots, m$. Thus $f_* p$ is saturated whenever

$$\rho_1(c) \oplus \dots \oplus \rho_r(c)(u_1), \dots, \rho_1(c) \oplus \dots \oplus \rho_r(c)(u_{m-1})$$

generate $M_{d_1} \oplus \dots \oplus M_{d_r}$ as a k -algebra. This is an open condition on c_{ij} . ■

5. PROOF OF THEOREM 1.2

In this section we complete the proof of Theorem 1.2. In view of Proposition 3.3, we only need to prove the following proposition.

PROPOSITION 5.1. *Assume $m \geq n + 1$. Then for any s -tuple (p_1, \dots, p_s) of distinct points of $Q_{m,n}$ there exists a $g \in \mathbf{Aut}(Q_{m,n})$ such that the s -tuple (gp_1, \dots, gp_s) is well behaved. Moreover, we can take $g = f_*$ for some $f \in \mathbf{TAut}(T_{m,n})$.*

To see how Theorem 1.2 follows from Proposition 3.3 and Proposition 5.1, let (p_1, \dots, p_s) and (q_1, \dots, q_s) be as in the statement of Theorem 1.2. By Proposition 5.1 we can find g and $\tilde{g} \in \mathbf{Aut}(Q_{m,n})$ such that the s -tuples (gp_1, \dots, gp_s) and $(\tilde{g}q_1, \dots, \tilde{g}q_s)$ are both well behaved. Hence, by Proposition 3.3 there exists an $h \in \mathbf{Aut}(Q_{m,n})$ such that $hgp_i = \tilde{g}q_i$ for $i = 1, \dots, s$. In other words, $\tilde{g}^{-1}hg$ takes p_i to q_i for each i , as desired.

We now turn to the proof of Proposition 5.1.

REDUCTION 1. *We may assume without loss of generality that $s = 2$.*

Proof. Suppose for each $i \neq j$ there exists an $f_{ij} \in \mathbf{TAut}(T_{m,n})$ such that $(f_* p_i, f_* p_j)$ is well behaved. Since

$$\mathbf{TAut}(T_{m,n}) = \bigcup_d G_d,$$

we can find a single d such that G_d contains every f_{ij} . For this d let W be the set of all $f \in G_d$ such that $(f_* p_1, \dots, f_* p_s)$ is well behaved, and let W_{ij} be the set of all $f \in G_d$ such that $(f_* p_i, f_* p_j)$ is well behaved. Note that

$$W = \bigcap_{1 \leq i < j \leq s} W_{ij};$$

see Definition 3.2. Each W_{ij} is nonempty by our choice of d and is open in G_d by Lemma 4.3. Since G_d is irreducible (Lemma 4.2), this implies that W is nonempty. ■

From now on we shall assume that $s = 2$ and denote p_1 and p_2 by p and q respectively.

REDUCTION 2. *We may assume without loss of generality that p and q are saturated.*

Proof. It is enough to show that there exists an $f \in \mathrm{TAut}(T_{m,n})$ such that f_*p and f_*q are saturated; we can then replace (p, q) by (f_*p, f_*q) .

In fact, we only need to show that there exists an $f \in \mathrm{TAut}(T_{m,n})$ such that f_*p is saturated. Indeed, let W_p be the set of all f such that f_*p is saturated, and let W_q be the set of all f such that f_*q is saturated. We want to show that $W_p \cap W_q \neq \emptyset$. Indeed, if $W_p \neq \emptyset$ and $W_q \neq \emptyset$, then for d sufficiently large $W_p \cap G_d \neq \emptyset$ and $W_q \cap G_d \neq \emptyset$. By Lemma 4.4 both $W_p \cap G_d$ and $W_q \cap G_d$ are Zariski-open in G_d . Since G_d is irreducible (Lemma 4.2), their intersection is nonempty. Hence, $W_p \cap W_q \neq \emptyset$, as desired.

Now let ϕ be the reduced semisimple representation associated to p . By [14, 8.1, 8.3 and 3.4] there exists a tame automorphism σ of F_m such that $\phi \circ \sigma$ is saturated. This means that $(\sigma')_*p$ is saturated. ■

Note that the last argument makes use of the assumption $m \geq n + 1$. From now on we shall assume that p and q are saturated. Let the semisimple representations associated to p and q be

$$\rho = \rho_1^{a_r} \oplus \cdots \oplus \rho_r^{a_r} \quad \text{and} \quad \mu = \mu_1^{b_1} \oplus \cdots \oplus \mu_s^{b_s}$$

respectively. Here $\rho_i, \mu_i : F_m \rightarrow M_{d_i}(k)$ are irreducible representations; $\rho_i \neq \rho_j, \mu_i \neq \mu_j$ for $i \neq j$.

Our further arguments rely on the following lemma.

LEMMA 5.2. *Let $x, y \in Q_{m,n}$. Denote their associated semisimple representations by*

$$\phi = \phi_1^{a_r} \oplus \cdots \oplus \phi_r^{a_r} \quad \text{and} \quad \nu = \nu_1^{b_1} \oplus \cdots \oplus \nu_s^{b_s}.$$

Assume that there is a polynomial $P(u_1, \dots, u_{m-1}) \in F_{m-1}$ such that

$$\mathrm{tr} \phi(P) \neq \mathrm{tr} \nu(P).$$

*Then there exists a tame automorphism f of $T_{m,n}$ such that (f_*p, f_*q) is well behaved.*

Proof. Consider the lower triangular automorphism f of $T_{m,n}$ given by

$$\begin{aligned} X_1 &\rightarrow X_1, \\ &\vdots \\ X_{m-1} &\rightarrow X_{m-1}, \\ X_m &\rightarrow \text{tr } P(X_1, \dots, X_{m-1}) + cX_m. \end{aligned}$$

We claim that (f_*p, f_*q) is well behaved for all but finitely many values of $0 \neq c \in k$. Denote the semisimple representation associated to f_*p and f_*q by

$$\phi(c) = \phi_1(c)^{a_1} \oplus \dots \oplus \phi_r(c)^{a_r} \quad \text{and} \quad \nu(c) = \nu_1(c)^{b_1} \oplus \dots \oplus \nu_s(c)^{b_s}.$$

Note the $\phi_i(c)(u_j) = \phi_i(u_j)$ and $\nu_i(c)(u_j) = \nu_i(u_j)$ for every $j = 1 \dots, m-1$, but

$$\begin{aligned} \phi_i(c)(u_m) &= \text{tr } \phi(P) I_{d_i} + c\phi_i(u_m), \\ \nu_i(c)(u_m) &= \text{tr } \nu(P) I_{d_i} + c\nu_i(u_m), \end{aligned}$$

where I_{d_i} is the $d_i \times d_i$ identity matrix.

Choose $i = 1, \dots, r$ and $j = 1, \dots, s$. It is enough to show that $\phi_i(c) \neq \nu_j(c)$ for all but finitely many c . This follows from the fact that

$$\text{tr } \phi_i(c)(u_m) = \text{tr } \nu_i(c)(u_m) \tag{6}$$

for at most finitely many c . Indeed, both sides of (6) are polynomials in c . These polynomials are not identical, since equality does not hold when $c = 0$. ■

REDUCTION 3. *We may assume without loss of generality that the restrictions of ρ and μ to $F_{m-1} = k\{u_1, \dots, u_{m-1}\}$ coincide.*

Proof. Recall that ρ and μ are defined up to conjugation in $M_n(k)$. Denote the restrictions of ρ and μ to F_{m-1} by $\tilde{\rho}$ and $\tilde{\mu}$ respectively. Since p and q are saturated, these representations are semisimple. If $\tilde{\rho} = g\tilde{\mu}g^{-1}$ for some $g \in \text{GL}_n(k)$, then we are done after replacing μ by $g\mu g^{-1}$.

We will now prove Proposition 5.1 under the assumption that no such g exists. Since $\tilde{\rho}$ and $\tilde{\mu}$ are semisimple, there is a polynomial $P(u_1, \dots, u_{m-1})$ such that

$$\mathrm{tr} \rho(P) \neq \mathrm{tr} \mu(P).$$

The result now follows from Lemma 5.2. ■

We are now ready to finish the proof of Proposition 5.1. Since p and q are distinct points of $Q_{m,n}$ and we are assuming that ρ and μ coincide on F_{m-1} , we must have $\rho(u_m) \neq \mu(u_m)$. Since ρ and μ are saturated, $\rho(u_m)$ and $\mu(u_m)$ lie in the semisimple subalgebra $\rho(F_{m-1}) = \mu(F_{m-1})$ of $M_n(k)$. Since the trace form on this subalgebra is nondegenerate, there exists an element $A(u_1, \dots, u_{m-1})$ such that

$$\mathrm{tr} \rho(u_m A) \neq \mathrm{tr} \mu(u_m A). \quad (7)$$

Let σ be a lower triangular automorphism of $T_{m,n}$ given by

$$\begin{aligned} u_1 &\rightarrow u_1, \\ &\vdots \\ u_{m-1} &\rightarrow u_{m-1}, \\ u_m &\rightarrow A(u_1, \dots, u_{m-1}) + cu_m. \end{aligned}$$

Note that

$$\mathrm{tr} \rho \circ \sigma(u_m^2) \neq \mathrm{tr} \mu \circ \sigma(u_m^2). \quad (8)$$

Indeed, both sides of (8) are quadratic polynomials in c . The left-hand side is equal to

$$\mathrm{tr} \rho(A^2) + 2c \mathrm{tr} \rho(u_m A) + c^2 \mathrm{tr} \rho(u_m^2),$$

and the right-hand side is

$$\mathrm{tr} \mu(A^2) + 2c \mathrm{tr} \mu(u_m A) + c^2 \mathrm{tr} \mu(u_m^2).$$

By (7) these quadratic polynomials have unequal linear terms. Hence, the inequality (8) holds for all but finitely many c , as claimed. We fix one such $c \neq 0$.

We are now ready to apply Lemma 5.2. Our polynomial $P(u_1, \dots, u_{m-1})$ will be u_1^2 . In view of (8), all we need to do now is permute the variables u_1 and u_m . More precisely, let δ be the linear automorphism of F_m given by $\delta(u_1) = u_m$, $\delta(u_m) = u_1$, and $\delta(u_i) = u_i$ for $i = 2, \dots, m-1$. Let $x = (\sigma')_* \circ (\delta')_* p$ and $y = (\sigma')_* \circ (\delta')_* q$. The associated semisimple representations of these points are $\phi = \rho \circ \sigma \circ \delta$ and $\nu = \mu \circ \sigma \circ \delta$ respectively. We can now rewrite (8) as

$$\mathrm{tr} \phi(P) \neq \mathrm{tr} \nu(P).$$

By Lemma 5.2 there exists a tame automorphism f of $T_{m,n}$ such that $(f_* x, f_* y)$ is well behaved. In other words, $(f_* \circ (\sigma')_* \circ (\delta')_* p, f_* \circ (\sigma')_* \circ (\delta')_* q)$ is well behaved. This completes the proof of Proposition 5.1.

6. A TRIVIALITY CRITERION AND THE GURALNICK-MONTGOMERY THEOREM

In this section we prove the following proposition.

PROPOSITION 6.1. *Suppose that $Z \in T_{m,n}$ has the same determinant as X_i for some $i = 1, \dots, m$.*

- (a) *Let $n \geq 3$. Then $Z = \zeta X_i$.*
- (b) *Let $n = 2$. Then $Z = \zeta X_i$ or $Z = \zeta(\mathrm{tr} X_i - X_i)$.*

Here $\zeta \in k$ is an n th root of unity.

As an immediate corollary of part (a) we obtain the following necessary and sufficient condition for $f \in \mathrm{Aut}(T_{m,n})$ to be trivial.

COROLLARY 6.2. *Suppose f is a k -algebra homomorphism of $T_{m,n}$ where $n \geq 3$. Then $f = \mathrm{id}$ if and only if for every $i = 1, \dots, m$ the element $f(X_i)$ has the same trace and the same determinant as X_i .*

Proof. Let $Z_i = f(X_i)$. By Proposition 6.1(a) $\det Z_i = \det X_i$ implies $Z_i = \zeta X_i$. Since $\mathrm{tr} Z_i = \mathrm{tr} X_i$, we must have $\zeta = 1$. In other words, $Z_i = f(X_i) = X_i$ for every $i = 1, \dots, m$. By Corollary 2.9 this is only possible if $f = \mathrm{id}$. ■

In the course of proving Proposition 6.1 we shall repeatedly appeal to the following elementary lemma.

LEMMA 6.3. *Let R be a commutative k -algebra, and let*

$$A(s) = A_d s^d + A_{d-1} s^{d-1} + \cdots + A_0$$

and

$$B(s) = B_d s^d + B_{d-1} s^{d-1} + \cdots + B_0$$

for some $A_1, \dots, A_d, B_0, \dots, B_d \in M_n(R)$. Suppose $\det A(s) = \det B(s)$ for every $s \in k$. Then $\det A_d = \det B_d$.

Proof. Note that $\det A(s) = s^{jd} \det A_d + O(s^{jd-1})$ and $\det B(s) = s^{jd} \det B_d + O(s^{jd-1})$. Since the base field k is infinite, this implies $\det A_d = \det B_d$. ■

We now recall that the trace ring $T_{m,n}$ is multigraded by the degree in X_1, \dots, X_m ; see Section 2. In particular, a monomial $Z = X_1^{i_1} \cdots X_m^{i_m}$ is homogeneous of multidegree (i_1, \dots, i_m) , and so is $\text{tr } Z$. The degree of an element $W \in T_{m,n}$ with respect to X_i will be denoted by $\deg_i(W)$.

We now proceed with the proof of Proposition 6.1. We may assume without loss of generality that $i = 1$.

LEMMA 6.4. *Suppose $\det Z = \det X_1$. Then $\deg_i(Z) = 0$ for $i = 2, \dots, m$, i.e., Z is independent of X_2, \dots, X_m .*

Proof. Assume the contrary, say $\deg_2(Z) = d \geq 1$. Write $Z = Z_d + Z_{d-1} + \cdots + Z_0$, where Z_i is a homogeneous element of degree i with respect to X_2 and $Z_d \neq 0$. Let $C = (C_1, \dots, C_m)$ be an m -tuple of matrices from $M_n(k)$. Let

$$\begin{aligned} A(s) &= Z(C_1, sC_2, C_3, \dots, C_m) \\ &= s^d Z_d(C) + s^{d-1} Z_{d-1}(C) + \cdots + Z_0(C). \end{aligned}$$

Applying Lemma 6.2 to $A(s)$ and $B(s) \equiv C_1$, we see that $\det Z_d(C) = 0$. Since the last equality holds for every m -tuple C of matrices from $M_n(k)$, we have $\det Z_d = 0$. By Proposition 2.4, this implies $Z_d = 0$, contradicting our assumption. ■

LEMMA 6.5. *Suppose $\det Z = \det X_1$. Then $\deg_1(Z) \leq 1$.*

Proof. By Lemma 6.4 we may assume $\deg_i(Z) = 0$ for $i = 2, \dots, m$, i.e., Z is independent of X_2, \dots, X_m . Let $\deg_1(Z) = d$, i.e., $Z = Z_d + Z_{d-1}$

$+ \cdots + Z_1 + Z_0$, where Z_i is a homogeneous element of degree i , $Z_d \neq 0$, and $d = \deg_1(Z)$. Substituting sA for X_1 , we see that sA has the same determinant as

$$Z(sA) = s^d Z_d(A) + s^{d-1} Z_{d-1}(A) + \cdots + Z_0(A).$$

We want to show $d \leq 1$. Assume the contrary. Then Lemma 6.2 with $A_i = Z_i(A)$, $B_1 = A$, and $B_2 = \cdots = B_d = 0$ says that $\det Z_d(A) = 0$. Moreover, this is true for every A , i.e., $\det Z_d = 0$. By Proposition 2.4, $Z_d = 0$, a contradiction. ■

We have therefore established that $Z = \alpha \operatorname{tr} X_1 + \beta X_1 + \gamma$ for some α , β , and $\gamma \in k$. Substituting the zero matrix for X_1 , we obtain $\gamma^n = \det 0 = 0$, i.e. $\gamma = 0$. Thus $Z = \alpha \operatorname{tr} X_1 + \beta X_1$. Substituting the diagonal matrix $\operatorname{diag}(\lambda_1, \dots, \lambda_n)$ for X_1 , we obtain a diagonal matrix

$$\operatorname{diag}(\alpha S + \beta \lambda_1, \dots, \alpha S + \beta \lambda_n)$$

whose determinant equals $\det X_1$. Here $\lambda_1, \dots, \lambda_n$ are independent commuting variables and $S = \lambda_1 + \cdots + \lambda_n$. In other words,

$$\prod_{i=1}^n (\alpha S + \beta \lambda_i) = \prod_{i=1}^n \lambda_i.$$

By unique factorization in $k[\lambda_1, \dots, \lambda_n]$ there are scalars $a_1, \dots, a_n \in k$ and a permutation $\sigma \in S_n$ such that $\alpha S + \beta \lambda_i = a_i \lambda_{\sigma(i)}$ for every $i = 1, \dots, n$. We can now finish the proof of Proposition 6.1.

(a) $n \geq 3$: The equality $\alpha S = -\beta \lambda_i + a_i \lambda_{\sigma(i)}$ is impossible unless $\alpha = 0$; just set $\lambda_i = \lambda_{\sigma(i)} = 0$ to get a contradiction. On the other hand, $\alpha = 0$ implies $Z = \beta X_1$. Taking the determinant, we see that $\beta^n = 1$.

(b) $n = 2$: In this case there are only two possibilities for the permutation σ : either $\sigma = \operatorname{id}$ or $\sigma = (1, 2)$.

Suppose $\sigma = \operatorname{id}$. Taking $i = 1$, we obtain $\alpha \lambda_1 + \alpha \lambda_2 = -\beta \lambda_1 + a_1 \lambda_1$ and thus $\alpha = 0$, i.e. $Z = \beta X_1$. Taking the determinant, we obtain $\beta^2 = 1$.

On the other hand, if $\sigma = (1, 2)$, then $\alpha \lambda_1 + \alpha \lambda_2 = -\beta \lambda_1 + a_1 \lambda_2$ and thus $\alpha = -\beta$. Hence, $Z = \alpha(\operatorname{tr} X_1 - X_1)$. Taking the determinant, we obtain $\alpha^n = 1$, and the proof is complete.

We now derive the following theorem of Guralnick and Montgomery from Proposition 6.1. (An analogous theorem for $G_{m,n}$ was proved by Montgomery in [9]; see also Lvov and Kharchenko [8] and Le Bruyn [7, Theorem 13].)

COROLLARY 6.6 (Guralnick and Montgomery [5]). *Let f be a k -algebra automorphism of the trace ring $T_{m,n}$. Suppose $f(c) = c$ for every c of $C_{m,n}$. Then*

- (a) *f is a trivial automorphism of $T_{m,n}$ if $n \geq 3$ or $m \geq 3$;*
- (b) *f is either trivial or is given by $f(X_i) = \text{tr } X_i - X_i$ ($i = 1, 2$) if $n = m = 2$.*

Proof. By Proposition 2.8

$$\det f(Z) = \det Z \quad \text{and} \quad \text{tr } f(Z) = \text{tr } Z \quad (9)$$

for every $Z \in T_{m,n}$.

(a): If $n \geq 3$, (9) implies $f = \text{id}$ by Corollary 6.2.

(b): If $n = 2$, then Proposition 6.1(b) implies $f(X_i) = X_i$ or $\text{tr } X_i - X_i$ for each $i = 1, \dots, m$. We claim that if the first possibility occurs for one i , then it has to occur for all other i . Indeed, suppose $f(X_1) = X_1$ but $f(X_2) = \text{tr } X_2 - X_2$. Then

$$\text{tr } X_1 X_2 = f(\text{tr } X_1 X_2) = \text{tr } [f(X_1)f(X_2)] = \text{tr } X_1 \text{tr } X_2 - \text{tr } X_1 X_2.$$

Substituting the elementary matrices e_{11} for X_1 and e_{22} for X_2 , we obtain $0 = 1$, a contradiction.

If $f(X_i) = X_i$ for some $i = 1, \dots, m$, then $f(X_i) = X_i$ for every i , and f is trivial by Corollary 2.9. The only other possibility is $f(X_i) = \text{tr } X_i - X_i$ for every $i = 1, \dots, m$. However, if such an automorphism existed for $m \geq 3$, then we would have

$$\text{tr } X_1 X_2 X_3 = \text{tr } f(X_1 X_2 X_3) = \text{tr } [(\text{tr } X_1 - X_1)(\text{tr } X_2 - X_2)(\text{tr } X_3 - X_3)].$$

Substituting the elementary matrices e_{12} for X_1 , e_{21} for X_2 , and e_{11} for X_3 , we obtain $1 = 0$, a contradiction. ■

7. INDUCED AUTOMORPHISMS

We now return to the group homomorphisms

$$\text{Aut}(F_m) \rightarrow \text{Aut}(T_{m,n}) \rightarrow \mathbf{Aut}(Q_{m,n}) \quad (10)$$

defined in Section 3. Recall that the first map is given by $\sigma \rightarrow \sigma'$ and the second map is given by $f \rightarrow f_*$.

THEOREM 7.1. *Let $m, n \geq 2$. In the diagram (10),*

- (a) *the second map is injective iff $(m, n) \neq (2, 2)$,*
- (b) *the first map is not surjective, and*
- (c) *the first map is injective iff $m = 2$.*

Proof. (a) is a direct consequence of the Guralnick-Montgomery theorem 6.6.

(b): By Proposition 2.10 there is an automorphism f of $T_{m,n}$ given by

$$\begin{aligned} X_1 &\rightarrow X_1 + \text{tr } X_2, \\ X_2 &\rightarrow X_2, \\ &\vdots \\ X_m &\rightarrow X_m. \end{aligned}$$

We claim that this automorphism is not induced by an automorphism of the polynomial ring F_m . Indeed, assume the contrary. Then we have an identity of the form $\text{tr } X_1 = P(X_1, \dots, X_m)$ for some polynomial $P \in F_m$. Since the left-hand side is multihomogeneous of degree $(1, 0, \dots, 0)$, the right-hand side must be equal to αX_1 for some $\alpha \in k$, a contradiction.

(c): We first consider the case $m \geq 3$. Let $r \in k\{u_1, u_2\}$ be a two-variable polynomial identity for the matrix algebra $M_n(k)$. For example, $r(u_1, u_2)$ can be defined as $S_{n+1}(u_1, u_1 u_2, \dots, u_1^n u_2)$, where S_{n+1} is the standard identity of degree $n + 1$. Now let $\sigma \in \text{Aut}(F_m)$ be given by

$$\begin{aligned} u_1 &\rightarrow u_1, \\ u_2 &\rightarrow u_2, \\ &\vdots \\ u_m &\rightarrow u_m + (u_1, u_2). \end{aligned}$$

Then $\sigma \neq \text{id}$ in $\text{Aut}(F_m)$ but $\sigma' = \text{id}$ in $\text{Aut}(T_{m,n})$, as desired.

Now suppose $m = 2$. Let $I_j \subset F_2$ be the ideal of polynomial identities of $j \times j$ matrices. Identifying F_m/I_n with $G_{m,n}$ and F_m/I_1 with the commutative polynomial ring $k[x_1, x_2]$, we see that the “abelianization” homomorphism $\text{Aut}(F_2) \rightarrow \text{Aut}(k[x_1, x_2])$ factors through $\text{Aut}(G_{2,n})$. The “abelianization” homomorphism is known to be an isomorphism in the two-variable case;

see [3]. Thus the map $\text{Aut}(F_2) \rightarrow \text{Aut}(G_{2,n})$ is injective. By Corollary 2.9 the map $\text{Aut}(F_2) \rightarrow \text{Aut}(T_{2,n})$ is also injective. ■

We now turn to the proof of Theorem 1.3, which says that the homomorphism $\text{Aut}(T_{m,n}) \rightarrow \mathbf{Aut}(Q_{m,n})$ in (10) is not surjective.

Let t be the automorphism of $[M_n(k)]^m$ (as an algebraic variety) given by

$$t : (A_1, \dots, A_m) \rightarrow (A_1^t, \dots, A_m^t).$$

Here A_i^t is the transpose of the $n \times n$ matrix A_i . Note that for any $g \in \text{PGL}_n$ and any $A = (A_1, \dots, A_m) \in (M_n)^m$ we have $t(gA) = (g^{-1})'(tA)$. Here gA stands for the m -tuple $(gAg^{-1}, \dots, gA_mg^{-1})$. By the universal property of the affine quotient $Q_{m,n} = [M_n(k)]^m / \text{PGL}_n$, t descends to an automorphism $t_0 : Q_{m,n} \rightarrow Q_{m,n}$. It easily follows from the definition of the representation type that $\tau(t_0(p)) = \tau(p)$ for every $p \in Q_{m,n}$. In other words, $t \in \mathbf{Aut}(Q_{m,n})$. We can now state and prove a more precise form of Theorem 1.3.

PROPOSITION 7.2. *Assume that $n \geq 3$ and $m \geq 2$. Then t_0 is not induced by an automorphism of $T_{m,n}$.*

Proof. Assume the contrary: $t_0 = f_*$ for some $f \in \text{Aut}(T_{m,n})$. Then by Proposition 2.8 we have

$$c_j(f(X_i)) = t_0(c_j(X_i)) = c_j(X_i^t) = c_j(X_i)$$

for every $i = 1, \dots, m$. Here $c_j(M)$ is the j th coefficient of the characteristic polynomial of M . In particular, $f(X_i)$ has the same trace and determinant as X_i for every $i = 1, \dots, m$. Since $n \geq 3$, Corollary 6.2 says that this is only possible when $f = \text{id}$.

Thus we only need to show that $t_0 \neq \text{id}$. Assume the contrary: $t_0 = \text{id}$. Then we must have

$$\text{tr}(X_1 X_2 X_1^2 X_2^2) = \text{tr}\left[X_1^t X_2^t (X_1^t)^2 (X_2^t)^2\right] = \text{tr}(X_2^2 X_1^2 X_2 X_1). \quad (11)$$

We now substitute $e_{12} + e_{23}$ for X_1 and $e_{21} + ae_{32}$ for X_2 into (11). Here e_{ij} is the (i, j) th elementary matrix in $M_n(k)$ and a is an element of k . The identity (11) then yields $a = a^2$ for every $a \in k$, a contradiction. ■

REMARKS.

(1) The last argument can be made more transparent if $m \geq 3$. Indeed, in this case $t_0 = \text{id}$ implies

$$\text{tr } X_1 X_2 X_3 = \text{tr } X_1' X_2' X_3' = \text{tr } X_3 X_2 X_1.$$

Substituting the elementary matrices e_{12} , e_{23} , and e_{13} for X_1 , X_2 , and X_3 respectively, we obtain $0 = 1$, a contradiction.

(2) For $n = 2$, t_0 is in fact induced by an automorphism of the trace ring. Let $Y_i = \text{tr } X_i - X_i$ for $i = 1, \dots, m$. Note that $X_i = \text{tr } Y_i - Y_i$. By Proposition 2.10 there exists an automorphism f of $T_{m,2}$ which takes X_i to Y_i . I claim that $t_0 = f_*$. Indeed, it is enough to check

$$\text{tr } P(fX_1, \dots, fX_m) = \text{tr } P(X_1^t, \dots, X_m^t)$$

for every polynomial $P \in F_m$. This is a consequence of the identity $M^t = g(\text{tr } M - M)g^{-1}$, where $g = e_{21} - e_{12}$ and M is an arbitrary 2×2 matrix.

The automorphism $t_0 = f_* : Q_{m,2} \rightarrow Q_{m,2}$ is clearly trivial when $m = 1$. (If $m = 1$ then $t = \text{id}$ and thus $t_0 = \text{id}$ for any n .) By the Guralnick-Montgomery theorem 6.6 it is also trivial when $m = 2$ and nontrivial for all other values of m .

I would like to thank Nikolaus Vonessen for helpful comments and suggestions.

REFERENCES

- 1 M. Artin, On Azumaya algebras and finite dimensional representations of rings, *J. Algebra* 11:532–563 (1969).
- 2 P. M. Cohn, *Algebra*, Vol. 2, 2nd ed., Wiley, New York, 1982.
- 3 A. J. Czerniakiewicz, Automorphisms of a free polynomial algebra of rank 2. II, *Trans. Amer. Math. Soc.* 171:309–315 (1972).
- 4 E. Formanek, Noncommutative invariant theory, in *Group Actions on Rings*, Contemp. Math. 13, 1985.
- 5 R. Guralnick and S. Montgomery, Normalizing elements of the trace ring of generic matrices, *Comm. Algebra* 17:1805–1813 (1989).
- 6 L. Le Bruyn, Simultaneous equivalence of square matrices, in *Seminaire d'Algebre Paul Dubreil et Marie-Paul Malliavin*, Lecture Notes in Math. 1404, Springer-Verlag, Berlin, 1989.
- 7 L. Le Bruyn, The Artin-Schofield theorem and some applications, *Comm. Algebra* 14:1439–1455 (1986).

- 8 I. V. L'vov and V. K. Kharchenko, Normal elements of the algebra of generic matrices are central (in Russian), *Sibirsk. Mat. Zh.* 23:193–195 (1982).
- 9 S. Montgomery, X -inner automorphisms of filtered algebras II, *Proc. Amer. Math. Soc.* 87 (1983).
- 10 D. Mumford and J. Fogarty, *Geometric Invariant Theory*, Springer-Verlag, New York, 1982.
- 11 C. Procesi, The invariant theory of $n \times n$ matrices, *Adv. Math.* 19:306–381 (1976).
- 12 C. Procesi, A formal inverse to the Cayley-Hamilton theorem, *J. Algebra* 107:63–74 (1987), 122:41–85 (1971).
- 13 Z. Reichstein, A functional interpretation of the ring of matrix invariants, *J. Algebra* 136:439–462 (1991).
- 14 Z. Reichstein, On automorphisms of matrix invariants, *Trans. Amer. Math. Soc.*, to appear.
- 15 L. H. Rowen, *Polynomial Identities in Ring Theory*, Academic, New York, 1980.
- 16 K. S. Sibirskii, Algebraic invariants of a set of matrices (in Russian), *Sibirsk. Mat. Zh.* 9(1):152–164 (1968); English transl., *Siberian Math. J.* 9:115–124 (1968).
- 17 I. R. Shafarevich, On some infinite-dimensional groups (in Russian), *Izv. Akad. Nauk SSSR Ser. Mat.* 45(1):214–226 (1981); English transl., *Math. USSR Izv.* 18(1):185–194 (1982).

Received 30 August 1992; final manuscript accepted 20 October 1992